

## Point groups and their representations

## Exercise 2.1 Energy bands of almost free electrons on the fcc lattice

Let us consider almost free electrons on a face-centered cubic (fcc) lattice. The goal of this exercise is to compute the lowest energy bands along the  $\Delta$ -line using degenerate perturbation theory and the machinery of *group theory*. Remember that in reciprocal space, the fcc lattice transforms into a body-centered cubic (bcc) lattice. The point group of the cubic Bravais lattices (simple cubic, fcc, bcc) is denoted by  $O_h$  (symmetry group of a cube). Its character table is given in Tab. 1.

- a) We first study the  $\Gamma$  point ( $\vec{k} = 0$ ). For *free* electrons ( $V = 0$ ) the lowest energy level is non-degenerate and the second one has an eight fold degeneracy. We focus on the second level and denote the eight-dimensional representation of  $O_h$  defined on this subspace by  $\Gamma$ . Find the irreducible representations contained in  $\Gamma$ . Compute the group character  $\chi_\Gamma$  and use the character table of  $O_h$  to show that

$$\Gamma = \Gamma_1^+ \oplus \Gamma_2^- \oplus \Gamma_4^- \oplus \Gamma_5^+ . \quad (1)$$

$O_h$	$E$ [ $xyz$ ]	$C_3(8)$ [ $zxy$ ]	$C_4^2(3)$ [ $\bar{x}\bar{y}\bar{z}$ ]	$C_2(6)$ [ $yx\bar{z}$ ]	$C_4(6)$ [ $\bar{y}xz$ ]	$J$ [ $\bar{x}\bar{y}\bar{z}$ ]	$JC_3(8)$ [ $\bar{z}\bar{x}\bar{y}$ ]	$JC_4^2(3)$ [ $xy\bar{z}$ ]	$JC_2(6)$ [ $\bar{y}\bar{x}\bar{z}$ ]	$JC_4(6)$ [ $y\bar{x}\bar{z}$ ]
$\chi_{\Gamma_1^+}$	1	1	1	1	1	1	1	1	1	1
$\chi_{\Gamma_1^-}$	1	1	1	1	1	-1	-1	-1	-1	-1
$\chi_{\Gamma_2^+}$	1	1	1	-1	-1	1	1	1	-1	-1
$\chi_{\Gamma_2^-}$	1	1	1	-1	-1	-1	-1	-1	1	1
$\chi_{\Gamma_3^+}$	2	-1	2	0	0	2	-1	2	0	0
$\chi_{\Gamma_3^-}$	2	-1	2	0	0	2	1	-2	0	0
$\chi_{\Gamma_4^+}$	3	0	-1	-1	1	3	0	-1	-1	1
$\chi_{\Gamma_4^-}$	3	0	-1	-1	1	-3	0	1	1	-1
$\chi_{\Gamma_5^+}$	3	0	-1	1	-1	3	0	-1	1	-1
$\chi_{\Gamma_5^-}$	3	0	-1	1	-1	-3	0	1	-1	1

Table 1: The character table of the cubic point group  $O_h$ .

- b) A finite periodic potential will in general split the second energy level at the  $\Gamma$  point. Applying degenerate perturbation theory to the Bloch equation [Eq. (1.20) in the lecture notes] leads to a  $8 \times 8$  matrix with off-diagonal elements  $u = V_{\frac{4\pi}{a}(1,1,1)}$ ,  $v = V_{\frac{4\pi}{a}(1,0,0)}$  and  $w = V_{\frac{4\pi}{a}(1,1,0)}$  (we basically follow chapter 1.3 of the lecture notes). This matrix can be diagonalized by going into the symmetry subspaces.

Show that for the energies and the wave functions one finds

$$\begin{aligned}
 \Gamma_1^+ &: E_0 + u + 3v + 3w && \cos\left(\frac{2\pi}{a}x\right) \cos\left(\frac{2\pi}{a}y\right) \cos\left(\frac{2\pi}{a}z\right); \\
 \Gamma_2^- &: E_0 - u - 3v + 3w && \sin\left(\frac{2\pi}{a}x\right) \sin\left(\frac{2\pi}{a}y\right) \sin\left(\frac{2\pi}{a}z\right); \\
 \Gamma_4^- &: E_0 - u + v - w && \left\{ \sin\left(\frac{2\pi}{a}x\right) \cos\left(\frac{2\pi}{a}y\right) \cos\left(\frac{2\pi}{a}z\right), \text{cyclic} \right\}; \\
 \Gamma_5^+ &: E_0 + u - v - w && \left\{ \cos\left(\frac{2\pi}{a}x\right) \sin\left(\frac{2\pi}{a}y\right) \sin\left(\frac{2\pi}{a}z\right), \text{cyclic} \right\};
 \end{aligned} \tag{2}$$

where  $E_0 = \frac{\hbar^2}{2m} 3\left(\frac{2\pi}{a}\right)^2$ .

- c) How do the irreducible representations split on the  $\Delta$ -line? The  $\Delta$ -line is defined by the points  $\vec{k} = \frac{\pi}{a}(0, 0, \delta)$ ,  $0 \leq \delta \leq 1$ . Use the character table of  $C_{4v}$ .

$C_{4v}$	$E$	$C_2(1)$	$C_4(2)$	$\sigma_v(2)$	$\sigma_d(2)$
	$[xyz]$	$[\bar{x}\bar{y}z]$	$[y\bar{x}z]$	$[\bar{x}yz]$	$[y\bar{x}z]$
$\chi_{\Delta_1}$	1	1	1	1	1
$\chi_{\Delta_2}$	1	1	1	-1	-1
$\chi_{\Delta_3}$	1	1	-1	1	-1
$\chi_{\Delta_4}$	1	1	-1	-1	1
$\chi_{\Delta_5}$	2	-2	0	0	0

Table 2: The character table of  $C_{4v}$ .

- d) Let us now consider the point  $X = \frac{2\pi}{a}(0, 0, 1)$ . The lowest level is two fold and the second four fold degenerate for  $V = 0$ . Compute the energies and the wave functions for these two levels.
- e) Finally, sketch the energy bands between the  $\Gamma$  and the  $X$  point. For an actual numerical calculation use the values  $u = -0.05$ ,  $v = 0.05$  and  $w = 0.1$  (in units of  $\frac{(2\pi\hbar)^2}{2ma^2}$ ).

### Exercise 2.2 Lifting the degeneracy of the atomic states

Determine how the energy levels of the  $p$ ,  $d$  and  $f$  orbitals of an atom lift due to a crystal field with cubic symmetry. Compute the corresponding eigenstates for the  $d$  orbitals. For this, consider the homogeneous harmonic polynomials of order 2. Alternatively, have a look at the basis functions given on page 11 of the lecture notes.