## Statistical Physics Exercise Sheet 8

HS 09 Prof. M. Sigrist

## Exercise 8.1 Independent Dimers in a Magnetic Field

We consider a system of N independent dimers of two spins, s=1/2, described by the Hamiltonian

$$\mathcal{H}_0 = J \sum_{i} (\vec{S}_{i,1} \cdot \vec{S}_{i,2}) \tag{1}$$

where i numbers the dimers and the second index m = 1, 2 denotes their magnetic site.

- a) What are the eigenstates and the eigenenergies of a single dimer? Consider the macroscopic system and determine the Helmholtz free energy, entropy, internal energy and specific heat as a function of temperature and N. Discuss the limit  $T \to 0$  and  $T \to \infty$  for both signs of J.
- b) We now apply a magnetic field in z direction leading to an additional term in the Hamiltonian,

$$\mathcal{H}' = -g\mu_B H \sum_{i,m} S_{i,m}^z. \tag{2}$$

How do the eigenenergies change? Sketch the energies with respect to the applied field H and determine the ground state. Discuss in this context the entropy per dimer in the limit  $T \to 0$ .

c) Calculate the magnetization m and the magnetic susceptibility  $\chi$  and discuss their dependence on H for different temperatures.

## Exercise 8.2 Spin Susceptibility of the Free Electron Gas

We consider a Fermi-gas with a Zeeman-coupling q to an external magnetic field B,

$$\mathcal{H}_B = \frac{g}{2}B(N_+ - N_-)\,, (3)$$

where  $N_{+}$  and  $N_{-}$  are the occupation numbers of the two spin species.

- 1. Derive a representation of the grand-canonical potential  $\Omega$  similar to the one in the lecture notes in the presence of this magnetic field. Show that it can be written as a sum  $\Omega = \sum_{\alpha} \Omega_{\alpha}$  over the spin species  $\alpha = \{+, -\}$  with a spin-dependent fugacity  $z_{\alpha}$ . Find an expression for the spin-susceptibility in terms of  $f_{n/2}(z_{\alpha})$  (for some n) by acting with an appropriate differential operator on  $\Omega$ .
- 2. Use the high-temperature expansion of  $f_{n/2}$  to calculate the corresponding limiting behavior of the spin-susceptibility in the limit  $B \to 0$ .
- 3. Use the low-temperature expansion to do the same for  $k_BT \ll \mu$ .

Office Hours: Monday, 9.11., 8 – 10 at HIT K 12.2.